

# Hip Evaluation Report

Report Date: 9/23/2011

Reference #: **896400**  
Practice #:

Radiography Date: 9/8/2011  
Date Received: 9/23/2011

**Owner:**  
ERIC GRAMLICH  
7158 STATE RD. U U  
FULTON, MO 65251  
UNITED STATES

**PennHIP Member:**  
DR. JENNIFER SHOCKLEY  
HIGGINSVILLE ANIMAL CLINIC  
2400 HWY BLVD  
HIGGINSVILLE, MO 64037  
UNITED STATES

ANIMAL	
OAK VIEW SUGAR	Reg. #: 2010001962011
CANINE / SOUTH AFRICAN BOERBOEL MASTIFF	Microchip: 0A01434471
Date of Birth: 7/21/2010 Sex: F Weight: 98 lbs. Age: 14 mo.	Tattoo:

RESULTS			
LEFT	Distraction Index (DI)	<b>0.54</b>	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	<b>None</b>	
	Cavitation	<b>No</b>	
	Other Findings	<b>Not Applicable</b>	
RIGHT	Distraction Index (DI)	<b>0.52</b>	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	<b>None</b>	
	Cavitation	<b>No</b>	
	Other Findings	<b>Not Applicable</b>	

Please note that the PennHIP DI is a measure of hip joint laxity, it does not allude to a "passing" or "failing" hip score.

LAXITY PROFILE RANKING										
The laxity profile ranking is based on the hip with the greater laxity (DI). This interpretation is based on a cross-section of 933 CANINE animals of the SOUTH AFRICAN BOERBOEL MASTIFF breed. The median DI for this group is 0.54.										
<b>Percentiles</b>										
	90th	80th	70th	60th	50th	40th	30th	20th	10th	
> 90th					Median					< 10th
↑										
The chart above indicates the ranking of your animal's passive hip laxity (DI) in relation to all CANINE animals of the SOUTH AFRICAN BOERBOEL MASTIFF breed in our database. Your animal's hip laxity lies within the 50th percentile or median range. Breed-specific evaluations are analyzed semi-annually. Consequently, the average laxity and range of laxity for any given group will change over time.										

PennHIP does not make specific breeding recommendations. Selection of sire and dam for mating is the decision of the breeder.

**NOTE: As a minimum breeding criterion, we propose that breeding stock be selected from the population of animals having hip laxity in the tighter half of the breed (to the left of the median mark on the graph). Higher selection pressure equates to more rapid expected genetic change per generation.**

By implementing selection based on passive hip laxity, we expect the breed average DI over the years to move toward tighter hip configuration, meaning lower hip dysplasia susceptibility. The PennHIP database permits scientific adjustment of criteria to reflect these shifts; the average laxity and range of laxity for a particular breed will change over time.